

References

- Athanasiadou, S. and Kyriazakis, I., 2004. Plant secondary metabolites: antiparasitic effects and their role in ruminant production systems. *Proceedings of the Nutrition Society*, 63(4), 631-639.
- Marley, C.L., Cook, R., Keatinge, R., Barrett, J. and Lampkin, N.H., 2003. The effect of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) on parasite intensities and performance of lambs naturally infected with helminth parasites. *Veterinary parasitology*, 112(1-2), 147-155.
- Marley, C.L., Cook, R., Barrett, J., Keatinge, R. and Lampkin, N.H., 2006. The effects of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) when compared with perennial ryegrass (*Lolium perenne*) on ovine gastrointestinal parasite development, survival and migration. *Veterinary Parasitology*, 138(3-4), 280-290.
- McFarland, C., Vineer, H.R., Chesney, L., Henry, N., Brown, C., Airs, P., Nicholson, C., Scollan, N., Lively, F., Kyriazakis, I. and Morgan, E.R., 2022. Tracking gastrointestinal nematode risk on cattle farms through pasture contamination mapping. *International Journal for Parasitology*, 52(10), 691-703.
- Molento, M.B., Buzatti, A. and Sprenger, L.K., 2016. Pasture larval count as a supporting method for parasite epidemiology, population dynamic and control in ruminants. *Livestock Science*, 192, 48-54.
- Rose, H., Wang, T., van Dijk, J. and Morgan, E.R., 2015. GLOWORM-FL: a simulation model of the effects of climate and climate change on the free-living stages of gastrointestinal nematode parasites of ruminants. *Ecological Modelling*, 297, 232-245.